

Vectree

Cardiovascular stasis and cellular preservation techniques in cryogenics

vectree.io/c/cellular-stasis-preservation

Cardiovascular stasis and cellular preservation techniques in cryogenics

vectree.io/c/cellular-stasis-preservation

Vascular Washout

Before cryopreservation can begin, the biological blood must be completely removed. Blood contains water that expands when frozen, and its components can clot or degrade, blocking the delivery of preservation fluids.

A surgical procedure involves cannulating the major vessels and using a pump to circulate a chilled base solution. This flushes out all erythrocytes and platelets, preparing the entire ‘plumbing’ system of the body for the introduction of chemical stabilizers.

Cryoprotectant Agents (CPAs)

Cryoprotectants are specialized chemical solutes (like Glycerol or DMSO) that act as ‘antifreeze’ for the body. Their primary role is to lower the freezing point of intracellular and extracellular fluids.

These agents penetrate the cell membranes and replace water molecules. By doing so, they disrupt the formation of hydrogen bonds required for ice crystals. However, CPAs are inherently toxic at high concentrations; the challenge lies in balancing concentration for protection against toxicity.

Vitrification

Vitrification is the ultimate goal of cellular preservation: the transformation of a liquid into a glass-like solid state without any crystal formation. In standard freezing, ice crystals act like microscopic jagged shards that shred cell membranes.

By using high concentrations of CPAs and rapid cooling, the molecules ‘freeze’ in place in a disordered, amorphous state. This prevents mechanical damage and locks the biological structures in time at temperatures below -130°C.

Thermal Gradient Management

Cooling is not a simple drop in temperature; it is a precisely managed descent. Computer-controlled cooling systems use liquid nitrogen vapor to lower the temperature at specific rates (e.g., 1°C per minute) to ensure uniform cooling across varying tissue densities.

Improper cooling rates can lead to ‘thermal fracturing,’ where different parts of the body shrink at different rates, causing macroscopic cracks in the tissue. Proper management ensures the transition to the cryogenic storage temperature of -196°C is structurally sound.

Cellular Integrity

At the microscopic level, preservation focuses on the lipid bilayer and the organelles. As the environment cools, the cell membrane becomes less fluid and more susceptible to rupture from osmotic pressure changes.

Techniques involve stabilizing the cytoskeleton and preventing the ‘out-flux’ of water that causes cell dehydration. If the cell’s internal structure can be maintained during the transition to a glass state, the biological information—including neurons and synaptic connections—remains theoretically intact.